

B. N. KHABIBULLIN

NEVANLINNA'S THEOREMS

Abstract. B. N. Khabibullin, *Nevanlinna's theorems*

We give a short survey on generalizations of Nevanlinna's theorems on zero distribution of bounded holomorphic functions and representation of meromorphic functions in multiply connected domains. It is a part of our report in the conference on complex analysis dedicated to the memory of Anatolii Asirovich Goldberg in Lviv, May 31-June 5, 2010.

Definitions, agreements, and basic notions. Let Ω be a domain in the complex plane $\mathbb{C}$ or in the Riemann sphere $\mathbb{C}_\infty \neq \Omega$. For $S \subset \Omega \subset \mathbb{C}_\infty$, we denote by $\overline{S}$ and $\partial\Omega$ the *closure* and the *boundary* of S relative to $\mathbb{C}_\infty$. We write $S \Subset \Omega$ if $\overline{S} \subset \Omega$.

Denote by $\text{Hol}(\Omega)$, $\text{Mer}(\Omega)$, $\text{sbh}(\Omega)$, and $\text{harm}(\Omega)$ the classes of all holomorphic, meromorphic, subharmonic, and harmonic functions on Ω .

We are concerned with finite or infinite sequences $\Lambda = \{\lambda_k\}$, $k = 1, 2, \dots$ of not necessarily distinct points from the domain Ω , without limit points in Ω . Let n_Λ be an integer-valued *counting measure* of sequence Λ defined by

$$n_\Lambda(S) := \sum_{\lambda_k \in S} 1, \quad S \subset \Omega.$$

Let $S \subset \Omega$. $\Lambda \subset S \iff \text{supp } n_\Lambda \subset S$.

The sequence Λ coincides with a sequence $\Gamma = \{\gamma_n\}$ (or is equal to Γ , or $\Lambda = \Gamma$) iff $n_\Lambda = n_\Gamma$. $\Lambda \subset \Gamma$ means $n_\Lambda \leq n_\Gamma$. $\Lambda \cap \Gamma$ and $\Lambda \cup \Gamma$ are defined by $n_{\Lambda \cap \Gamma} := \min\{n_\Lambda, n_\Gamma\}$ and $n_{\Lambda \cup \Gamma} := n_\Lambda + n_\Gamma$.

Given $f: A \rightarrow B$ and $b \in B$, we write $f \equiv b$ on A' if f is identically equal to b on $A' \subset A$; in the opposite case, $f \not\equiv b$ on A' .

Let $A, B \subset [-\infty, +\infty]$. A function $f: A \rightarrow B$ is *increasing* (*decreasing* resp.) if, for any $x_1, x_2 \in A$, $x_1 \leq x_2$ implies $f(x_1) \leq f(x_2)$ ($f(x_1) \geq f(x_2)$ resp.).

Given $a \in \mathbb{R}$, and $f: A \rightarrow [-\infty, +\infty]$, we set $a^+ := \max\{0, a\}$, $f^+ := \max\{0, f\}$.

The term “positive” (“negative” resp.) means “ ≥ 0 ” (“ ≤ 0 ” resp.).

Let $f \in \text{Hol}(\Omega)$ or $f \in \text{Mer}(\Omega)$, $f \not\equiv 0, \infty$ on Ω . Write Zero_f for the zero set of f (counting multiplicities). Evidently, Zero_f is a sequence of not necessarily distinct points from the domain Ω , without limit points in Ω .

A sequence Λ is the **zero sequence** for a subspace $H \subset \text{Hol}(\Omega)$ (further we write $\Lambda \in \text{Zero}(H)$) if and only if there exists a function $f \in H$ such that $\Lambda = \text{Zero}_f$.

A function $f \in \text{Hol}(\Omega)$ *vanish on* Λ if and only if $\Lambda \subset \text{Zero}_f$ (we write $f(\Lambda) = 0$).

A sequence Λ is the **zero subsequence** or the **non-uniqueness sequence** for the space H iff there exists a nonzero function $f \in H$ such that $f(\Lambda) = 0$.

Problems. Let H be a subspace of $\text{Hol}(\Omega)$. We consider the following five problems:

1. What point sequences Λ can be **zero sequences** for H ?
2. What point sequences Λ can be **zero subsequences** for H ?

3. In what cases is a zero **subsequence** for H simultaneously **zero sequence** for H or for some, preferably minimal, **extension space** $\widehat{H} \supset H$ of holomorphic functions in Ω ?
4. When can a **meromorphic function** in Ω be represented as a ratio of **holomorphic functions from H** ?
5. When can a **meromorphic function** in Ω be represented as a ratio of holomorphic functions from H **without common zeros**?

Green's function. Denote by

$$g_\Omega(\cdot, z): \mathbb{C}_\infty \setminus \{z\} \rightarrow [0, +\infty)$$

the *extended Green's function* for Ω with a pole at $z \in \Omega$, i.e., $g_\Omega(\zeta, z) \equiv 0$ for points $\zeta \in \mathbb{C}_\infty \setminus \overline{\Omega}$, $g_\Omega(\cdot, z) \in \text{sbh}(\mathbb{C}_\infty \setminus \{z\})$, $g_\Omega(\cdot, z) \in \text{harm}(\Omega \setminus \{z\})$, and also

$$g_\Omega(\zeta, z) = -\log|\zeta - z| + O(1), \quad \zeta \rightarrow z.$$

Given a continuous function $\phi: \partial\Omega \rightarrow \mathbb{R}$, we denote by $H_\Omega\phi$ the *solution the Dirichlet problem* for Ω with boundary function ϕ or the associated *Perron function*

$$H_\Omega\phi := \sup\{u \in \text{sbh}(\Omega) : \limsup_{z \rightarrow \zeta} u(z) \leq \phi(\zeta), \forall \zeta \in \partial\Omega\}.$$

Harmonic measure. Denote by $\mathcal{B}(\partial\Omega)$ the σ -algebra of Borel subset of $\partial\Omega$.

Denote by $\omega_\Omega(z, \cdot)$ the *harmonic measure* for Ω at the point $z \in \Omega$, i.e.,

$$\omega_\Omega(\cdot, \cdot): \Omega \times \mathcal{B}(\partial\Omega) \rightarrow [0, 1]$$

such that

- a) the map $B \mapsto \omega_\Omega(z, B)$ is a Borel probability measure on $\partial\Omega$;
- b) if $\phi: \partial\Omega \rightarrow \mathbb{R}$ is continuous function, then

$$H_\Omega\phi(z) = \int_{\partial\Omega} \phi(\zeta) d\omega_\Omega(z, \zeta).$$

The unit disk. Starting points of our researches are Nevalinna's theorems (1929).

Denote by $\text{Hol}^\infty(\mathbb{D}) \subset \text{Hol}(\mathbb{D})$ the space of holomorphic bounded functions on $\mathbb{D}$.

Denote by f_Λ a holomorphic function $\not\equiv 0$ on $\mathbb{D}$ with zero sequence $\text{Zero}_{f_\Lambda} = \Lambda \subset \Omega$.

Nevalinna's Theorem (on zeros). The following statements are equivalent:

- 1) Λ is a zero sequence for $\text{Hol}^\infty(\mathbb{D})$;
- 2) Λ is a zero subsequence for $\text{Hol}^\infty(\mathbb{D})$;
- 3) for each or some function $f_\Lambda \in \text{Hol}(\mathbb{D})$

$$\begin{aligned} \sup_{r < 1} (H_{r\mathbb{D}} \log |f_\Lambda|)(0) &= \sup_{r < 1} \int_{r\partial\mathbb{D}} \log |f_\Lambda(z)| d\omega_{r\mathbb{D}}(0, z) \\ &= \sup_{r < 1} \frac{1}{2\pi} \int_0^{2\pi} \log |f_\Lambda(re^{i\theta})| d\theta < +\infty; \end{aligned}$$

$$4) \sum_k g_{\mathbb{D}}(\lambda_k, 0) = \sum_k \log \frac{1}{|\lambda_k|} < +\infty \iff \sum_k (1 - |\lambda_k|) < +\infty.$$

Let $z \in \Omega$, $p, q \in \text{Hol}(\Omega)$, and let

$$f = \frac{p}{q} \in \text{Mer}(\Omega), \quad p(z) = q(z) = 1. \quad (1)$$

We set

$$u_f := \max\{\log |p|, \log |q|\} \in \text{sbh}(\Omega) \quad (2)$$

$$\text{or } u_f := \log \sqrt{|p|^2 + |q|^2} \in \text{sbh}(\Omega). \quad (3)$$

Let D be a subdomain of Ω , $z \in D \Subset \Omega$. The integral

$$T_D(f; z) := \int_{\partial D} u_f d\omega_D(z, \cdot) \quad (4)$$

is *Nevanlinna's characteristic* of f on D (in the form of Ahlfors–Shimizu for (3)) at the point z relative to the representation (1).

If $\text{Pol}_f = \{\gamma_k\}_{k=1}^\infty \subset \Omega$ is the *pole sequence* of f in Ω , i. e., zero sequence of $1/f$ in Ω , and $\text{Zero}_p \cap \text{Zero}_q = \emptyset$, then, for (2),

$$T_D(f; z) = \sum_k g_D(\gamma_k, z) + \int_{\partial D} \log^+ |f| d\omega_D(z, \cdot).$$

If $z = 0 \in D$, then $T_D(f) := T_D(f; 0)$.

Multiply connected domains. There are generalizations of Nevanlinna's theorems to classes of holomorphic and meromorphic functions on special finitely connected domains Ω .

Given $z \in \mathbb{C}$ and $0 < t < +\infty$, we write

$$D(z, t) := \{w \in \mathbb{C} : |w - z| < t\}, \quad \overline{D}(z, t) := \overline{D(z, t)}.$$

We consider now results from [1] (Andriy Kondratyuk, and Ilpo Laine, 2006). The authors develop the Nevanlinna theory and connected topics for meromorphic and holomorphic functions to $(m+1)$ -connected domains Ω , $m \in \mathbb{N}$, for following cases.

1. An $(m+1)$ -connected domain $\Omega = \mathbb{C} \setminus \bigcup_{j=1}^m \{c_j\}$, $c_j \in \mathbb{C}$, $m \in \mathbb{N}$, is called the *m-punctured plane*. For example, such 2-connected domain is the *punctured plane* $\mathbb{C} \setminus \{0\}$.
2. A bounded¹ $(m+1)$ -connected domain

$$\Omega = D(0, R) \setminus \left(\bigcup_{j=1}^m D(z_j, r_j) \right), \quad m \in \mathbb{N}, \quad D(z_j, t_j) \Subset D(0, R),$$

$\overline{D}(z_j, t_j) \cap \overline{D}(z_{j'}, t_{j'}) = \emptyset$ for all $j \neq j'$, is called *the strictly circular domain*. 2-connected annuli $A_R = D(0, R) \setminus D(0, 1/R)$, $1 < R < +\infty$, are examples of such domains.

¹always bounded relative to $\mathbb{C}$

3. Let $\Omega \subset \mathbb{C}$ be a bounded $(m+1)$ -connected domain, $m \in \mathbb{N}$,

$$\Omega = \bigcup_{j=0}^m G_j,$$

where G_j are simply connected domains (relative to $\mathbb{C}_\infty$), ∂G_j are simple (Jordan) closed paths, G_0 is bounded, and $\infty \in G_j$ for $j = 1, \dots, m$, $\mathbb{C} \setminus G_j \subseteq G_0$,

$$(\mathbb{C} \setminus G_j) \cap (\mathbb{C} \setminus G_{j'}) = \emptyset, \quad j \neq j', \quad j, j' \geq 1.$$

Such domains Ω are called in [1] admissible. Let φ_j are conformal mappings of $\mathbb{D}$ onto G_j realizing a conformal equivalence of $\mathbb{D}$ and G_j with $\varphi_0(0) = 0$, and $\varphi_j(0) = \infty$ as $j = 1, \dots, m$. Then every φ_j can be extended to a homeomorphism of $\overline{\mathbb{D}}$ onto $\overline{G_j}$ (the Carathéodory Theorem). For a simple closed path Γ , $\text{int } \Gamma$ denote the bounded domain with the boundary Γ (the Jordan Theorem).

Let

$$\Gamma_{jr}(s) := \varphi_j(e^{is}), \quad 0 \leq s \leq 2\pi, \quad j = 0, \dots, m, \quad \Gamma_{jr}^* := \Gamma_{jr}([0, 2\pi)), \quad j = 0, \dots, m,$$

and

$$\Omega_r := (\text{int } \Gamma_{0r}^*) \setminus \bigcup_{j=1}^k \overline{\text{int } \Gamma_{jr}^*}, \quad r_0 \leq r < 1.$$

Let $\Lambda = \{\lambda_k\}$ be a sequence in Ω , and let

$$N_0(r, \Lambda) := \int_{r_0}^r \frac{n_\Lambda(\Omega_t)}{t} dt, \quad r_0 \leq r < 1,$$

where $r_0 < 1$ is a constant sufficiently close to 1.

For a meromorphic function f on an admissible bounded domain Ω denote

$$m_0(r, f) := \frac{1}{2\pi} \sum_{j=1}^m \left(\int_0^{2\pi} \log^+ |f(\varphi_j(re^{is}))| ds - \int_0^{2\pi} \log^+ |f(\varphi_j(r_0 e^{is}))| ds \right),$$

$r_0 \leq r < 1$. The function

$$T_0(r, f) := N_0(r, \text{Pol}_f) + m_0(r, f), \quad r_0 \leq r < 1,$$

is the *Nevalinna characteristic* of f (in the sense of A. Kondratyuk and I. Laine).

Theorem K–L (on meromorphic functions, [1; Theorem 43.2]). *Let $f \in \text{Mer}(\Omega)$, where $\Omega \subset \mathbb{C}$ is a finitely connected bounded admissible domain. If $T_0(r, f) = O(1)$, $r \rightarrow 1$, then there are bounded functions $h_1, h_2 \in \text{Hol}(\Omega)$ such that $f = h_1/h_2$.*

Let $f_\Lambda \in \text{Hol}(\Omega)$ be a function with zero sequence $\Lambda \subset \Omega$. If we apply this theorem to the function $f_\Lambda \in \text{Hol}(\Omega)$, then we get

Theorem K–L (on zeros). *Let $\Omega \subset \mathbb{C}$ be a finitely connected admissible bounded domain. If*

$$\sup_{r_0 \leq r < 1} m_0(r, f_\Lambda) < +\infty,$$

then there is a bounded holomorphic on Ω function f such that $f(\Lambda) = 0$.

We consider the Problems 1–5 in a general form in [2], [3].

A function (weight) $M: \Omega \rightarrow \mathbb{R}$ define weighted space $\text{Hol}(\Omega; M) :=$

$$\left\{ f \in \text{Hol}(\Omega) : \sup_{z \in \Omega} \frac{|f(z)|}{\exp M(z)} < +\infty \right\}.$$

If $M \equiv 0$ on Ω , then $\text{Hol}(\Omega; 0) = \text{Hol}^\infty(\Omega)$.

Let Ω be a domain in $\mathbb{C}_\infty$.

We use following classification of domains Ω such that $\boxed{0 \in \Omega \not\rightarrow \infty}$.

I. This domain Ω is

- a) **simply connected** (for example, $\Omega = \mathbb{C}$ or $\Omega = \mathbb{D}$) or
- b) **finitely connected** and $\boxed{\overline{\Omega} \neq \mathbb{C}_\infty}$ (for example, each bounded finitely connected domain Ω).

II. This domain Ω is $(m+1)$ -**connected** with $m \in \mathbb{N}$ and $\boxed{\overline{\Omega} = \mathbb{C}_\infty}$.

III. This domain Ω is

- a) **bounded** or
- b) **unbounded**.

Let $\text{Exh } \Omega = \{D\}$ be a *exhaustion* of Ω , i. e., $\cup_{D \in \text{Exh } \Omega} D = \Omega$, where $0 \in D$ and every $D \in \text{Exh } \Omega$ is the regular domain for the Dirichlet problem. Such exhaustion exists always (countable increasing and such that all D have smooth boundary).

Theorem 1 (on zeros). *Let $\Lambda \subset \Omega$, and let Ω be a domain of type **I**. Following statements are equivalent:*

- 1) Λ is a zero sequence for $\text{Hol}^\infty(\Omega)$;
- 2) Λ is a zero subsequence for $\text{Hol}^\infty(\Omega)$;
- 3) for each or some function $f_\Lambda \in \text{Hol}(\Omega)$ with $\text{Zero}_{f_\Lambda} = \Lambda$

$$\sup_{D \in \text{Exh } \Omega} (H_D \log |f_\Lambda|)(0) := \sup_{D \in \text{Exh } \Omega} \int_D \log |f_\Lambda(z)| d\omega_D(0, z) < +\infty;$$

- 4) $\sup_{D \in \text{Exh } \Omega} \sum_k g_D(\lambda_k, 0) < +\infty$.

If Ω possesses the Green's function, then we can remove $\boxed{\sup_{D \in \text{Exh } \Omega}}$ everywhere and replace D with Ω .

Theorem 1 (on meromorphic functions). *Let Ω be a domain of type **I**. Let (see (1))*

$$f = \frac{p}{q} \in \text{Mer}(\Omega), \quad p, q \in \text{Hol}(\Omega), \quad p(0) = q(0) = 1,$$

and (see (4), (2), (3) resp.)

$$T_D(f) := \int_{\partial D} u_f d\omega_D(0, \cdot), \quad D \in \text{Exh}(\Omega),$$

where

$$u_f := \max\{\log |p|, \log |q|\} \in \text{sbh}(\Omega)$$

$$\text{or } u_f := \log \sqrt{|p|^2 + |q|^2} \in \text{sbh}(\Omega).$$

Assume that one of the following two conditions holds:

- 1) $\sup_{D \in \text{Exh } \Omega} T_D(f) < +\infty$;
- 2) $p, q \in \text{Hol}^\infty(\Omega)$.

Then there are $p_0, q_0 \in \text{Hol}^\infty(\Omega)$ **without common zeros** such that $f = p_0/q_0$.

Remark 1. If the domain Ω is regular for the Dirichlet problem, then we can remove

$$\sup_{D \in \text{Exh } \Omega}$$

in 1) and replace D with G .

Theorem 2 (on zeros). *Let $\Lambda \subset \Omega$, and let Ω be a domain of type **II**. If Λ is a zero sequence for $\text{Hol}^\infty(\Omega)$, then*

- 2) Λ is a zero subsequence for $\text{Hol}^\infty(\Omega)$;
- 3) for each (for some) function $f_\Lambda \in \text{Hol}(\Omega)$ with $\text{Zero}_{f_\Lambda} = \Lambda$

$$\sup_{D \in \text{Exh } \Omega} (H_D \log |f_\Lambda|)(0) := \sup_{D \in \text{Exh } \Omega} \int_D \log |f_\Lambda(z)| d\omega_D(0, z) < +\infty;$$

- 4) $\sup_{D \in \text{Exh } \Omega} \sum_k g_D(\lambda_k, 0) < +\infty$.

Conversely, assume that one of the conditions 2)–4) holds. Then there is a constant $b < m$ such that Λ is zero sequence for every space $\text{Hol}(\Omega, M)$ with

$$M: z \mapsto c_0^+ \log^+ |z| + \sum_{k=1}^m c_k^+ \log^+ \frac{1}{|z - a_k|}, \quad (5)$$

where $\sum_{k=0}^m c_k = b$, $a_k \in \mathbb{C} \setminus \Omega$.

Theorem 2 (on meromorphic functions). *Let Ω be a domain of type **II**. Let (see (1))*

$$f = \frac{p}{q} \in \text{Mer}(\Omega), \quad p, q \in \text{Hol}(\Omega), \quad p(0) = q(0) = 1.$$

Assume that one of the following two conditions holds:

- 1) $\sup_{D \in \text{Exh } \Omega} T_D(f) < +\infty$;
- 2) $p, q \in \text{Hol}^\infty(\Omega)$.

Then there are constant $b < m$ such that, for every function M from (5), there exist functions $p_0, q_0 \in \text{Hol}(\Omega; M)$ **without common zeros** representing $f = p_0/q_0$.

Theorem 3 (on zeros). *Let $\Omega \subset \mathbb{C}$ be a domain, and let $\Lambda \subset \Omega$ be a sequence. If Λ is a zero (sub)sequence for $\text{Hol}^\infty(\Omega)$, then*

3) for each or some function $f_\Lambda \in \text{Hol}(\Omega)$ with $\text{Zero}_{f_\Lambda} = \Lambda$

$$\sup_{D \in \text{Exh } \Omega} (H_D \log |f_\Lambda|)(0) := \sup_{D \in \text{Exh } \Omega} \int_D \log |f_\Lambda(z)| d\omega_D(0, z) < +\infty;$$

4) $\sup_{D \in \text{Exh } \Omega} \sum_k g_D(\lambda_k, 0) < +\infty$.

Conversely, let one of the conditions 3), 4) is fulfilled. Then Λ is zero subsequence for space $\text{Hol}(\Omega, M)$ with

$$M: z \mapsto \log \frac{1}{\text{dist}(z, \partial\Omega)}, \quad z \in \Omega,$$

where $\text{dist}(z, \partial\Omega)$ is the Euclidean distance from z up to $\partial\Omega$, if Ω is bounded, and with any

$$M: z \mapsto \log \frac{1}{\text{dist}(z, \partial\Omega)} + c_0 \log^+ |z| + c_1 \log^+ \frac{1}{|z - a|}, \quad z \in \Omega,$$

where $c_0 + c_1 = 9$, $a \in \mathbb{C} \setminus \Omega$, if Ω is unbounded.

Theorem 3 (on meromorphic functions). *Let Ω be a subdomain of $\mathbb{C}$. Let (see (1))*

$$f = \frac{p}{q} \in \text{Mer}(\Omega), \quad p, q \in \text{Hol}(\Omega), \quad p(0) = q(0) = 1.$$

Let's choose a function M as in the Theorem 3 (on zeros). Suppose that

$$\sup_{D \in \text{Exh } \Omega} T_D(f) < +\infty.$$

Then there exist functions $p_0, q_0 \in \text{Hol}(\Omega; M)$ representing $f = p_0/q_0$.

Remark 2. If Ω possesses the Green's function, then we can remove $\sup_{D \in \text{Exh } \Omega}$ everywhere in the Theorems 3 and replace D with Ω .

General results. If $M \in \text{sbh}(\Omega)$ with the Riesz measure $\nu_M := \frac{1}{2\pi} \Delta M \geq 0$, then there is always a global Riesz representation (decomposition)

$$M(z) = \int_{\Omega} k(\zeta, z) d\nu_M(\zeta) + H(z), \quad z \in \Omega, \quad (6)$$

where $H \in \text{harm}(\Omega)$,

$$k(\zeta, z) = \log |\zeta - z| + h_M(\zeta, z) \quad (7)$$

is a special subharmonic kernel with harmonic component $h(\zeta, z)$ of $z \in \Omega$ for each $\zeta \in \Omega$. Let $Q: \Omega \rightarrow \mathbb{R}$ be an upper semicontinuous function such that

$$\int_{\Omega} (k(\zeta, 0) - k(\zeta, z))^+ d\nu_M(\zeta) \leq Q(z) \quad (8)$$

for almost all $z \in \Omega$ in the Lebesgue measure on Ω .

Denote by $\mathcal{U}_0^d(\Omega)$ the class of all connected unions $D \ni 0$ of finitely many open disks from Ω whose complement has no one-point connected components.

Main Theorem (see [2, Theorems 1–5]). *Let $M \in \text{sbh}(\Omega) \cap C(\Omega)$ with the Riesz measure ν_M .*

[Z] *If $\Lambda = \{\lambda_k\} \subset \Omega$ is a zero (sub)set for $\text{Hol}(\Omega; M)$, then*

$$\sup_{0 \in D \in \Omega} \left(\sum_k g_D(\lambda_k, 0) - \int g_D(\zeta, 0) d\nu_M(\zeta) \right) < +\infty. \quad (9)$$

Conversely, if we have (9) where domains D run only the class $\mathcal{U}_0^d(\Omega)$, then Λ is zero subsequence (non-uniqueness sequence) for $\text{Hol}(\Omega; \widehat{M})$ where $\widehat{M}(z) :=$

$$\inf_{0 < t < \text{dist}(z, \partial\Omega)} \left(\frac{1}{2\pi} \int_0^{2\pi} M(z + te^{i\theta}) d\theta + \log(1 + 1/t) \right) + 9 \log^+ |z|,$$

and zero sequence for $\text{Hol}(\Omega; M + Q)$.

Thus, every zero subsequence for $\text{Hol}(\Omega; M)$ is zero sequence for $\text{Hol}(\Omega; M + Q)$.

[M] *Let $f = g/q$ be a meromorphic function and $g, q \in \text{Hol}(\Omega; M)$. Then*

$$\sup_{0 \in D \in \Omega} \left(\int_{\Omega} \log \max\{|g|, |q|\}(z) d\omega_D(0, z) - \int M(z) d\omega_D(0, z) \right) < +\infty. \quad (10)$$

Conversely, if, under the assumptions of (6) and (8), we have (10) where domains D run only the class $\mathcal{U}_0^d(\Omega)$, then there are functions $g, q \in \text{Hol}(\Omega; \widehat{M})$ and $g_0, q_0 \in \text{Hol}(\Omega; M + Q)$ such that $f = g/q = g_0/q_0$ and g_0, q_0 have not common zeros.

Thus, if $f = g/q$ with $g, q \in \text{Hol}(\Omega; M)$, then there exist functions $g_0, q_0 \in \text{Hol}(\Omega; M + Q)$ without common zeros such that $f = g_0/q_0$.

Nevanlinna's theorems wit nonradial and nonpositive weight. In [4, Theorem 1], we investigate also an slowly counterpart Nevanlinna's theorems.

Let $M: \mathbb{D} \rightarrow \mathbb{R}$, and let $M \in \text{sbh}(\mathbb{D})$ with the Riesz measure ν_M . Given $z = re^{i\theta}$, $0 \leq r < 1$, $\theta \in \mathbb{R}$, and $a > 0$ we consider a polar rectangle

$$\Xi(z; a) := \{ \zeta = te^{i\psi} : (r - a\sqrt{1-r^2})^+ \leq t < 1, |\sin(\psi - \theta)| < a\sqrt{1-r^2} \} \quad (11)$$

of relative size a and the function

$$q_M^{[a]}(z) := \frac{1}{1-|z|} \int_{\Xi(z; a)} (1-|\zeta|) d\nu_M(\zeta). \quad (12)$$

We set

$$A_M^{[\varepsilon]}(z) := \frac{1}{2\pi} \int_0^{2\pi} M(z + \varepsilon(1-|z|)e^{i\theta}) d\theta, \quad 0 < \varepsilon < 1. \quad (13)$$

Theorem 4. *Let M be a subharmonic function $\mathbb{D}$, $M(0) > -\infty$ and*

$$\sup_{r < 1} \int_0^{2\pi} M(re^{i\theta}) d\theta < +\infty, \quad (14)$$

that is equivalent a Blaschke condition

$$\int_0^1 (1-t) d\nu_M(t) < +\infty. \quad (15)$$

For a function f_Λ with $\text{Zero}_{f_\Lambda} = \Lambda$,

(Z) if is fulfilled at least one condition

$$\sup_{D \in \mathcal{U}_0^d(\mathbb{D})} \left(\int_{\mathbb{D}} \log |f_{\Lambda}(z)| d\omega_D(0, z) - \int_{\mathbb{D}} M(z) d\omega_D(0, z) \right) < +\infty, \quad (16f)$$

$$\sup_{D \in \mathcal{U}_0^d(\mathbb{D})} \left(\sum_k g_D(\lambda_k, 0) - \int_{\mathbb{D} \setminus \{0\}} g_D(\zeta, 0) d\nu_M(\zeta) \right) < +\infty, \quad (16g)$$

$$\sup_{D \in \mathcal{U}_0^d(\mathbb{D})} \left(\sum_k g_D(\lambda_k, 0) - \int_{\mathbb{D}} M(z) d\omega_D(0, z) \right) < +\infty, \quad (16h)$$

then for any $\varepsilon \in (0, 1)$ and $1 < a < 2$ the sequence $\Lambda \subset \mathbb{D}$ is zero sequence for space

$$\text{Hol}\left(\mathbb{D}; A_M^{[\varepsilon]} + \frac{C_\varepsilon}{2-a} q_M^{[a]}\right), \quad (17)$$

where a constant C_ε dependent only on ε ;

(U) if Λ is zero subsequence for $\text{Hol}(\mathbb{D}; M)$, then Λ is zero sequence for (17);

(M) if meromorphic function $f = g/q$ in $\mathbb{D}$ is represented as a ratio of functions $g, q \in \text{Hol}(\mathbb{D})$, $\max\{|g(0)|, |q(0)|\} \neq 0$, and is fulfilled at least one condition

$$\sup_{D \in \mathcal{U}_0^d(\mathbb{D})} \left(\int_{\mathbb{D}} \log \max\{|g(z)|, |q(z)|\} d\omega_D(0, z) - \int_{\mathbb{D}} M(z) d\omega_D(0, z) \right) < +\infty, \quad (18h)$$

$$g, q \in \text{Hol}(\mathbb{D}; M), \quad (18M)$$

then there are functions g_0 and q_0 from class (17) without common zero, such that $f = g_0/q_0$ in $\mathbb{D}$.

A function $M: z = re^{i\theta} \rightarrow [-\infty, +\infty]$ is called *radial in a sector*

$$\angle(\alpha, \beta) := \{z = re^{i\theta}: 0 \leq r < 1, \alpha < \theta < \beta\} \quad (19)$$

from $\mathbb{D}$, if for each $0 \leq r < 1$ the function $M(re^{i\theta})$ independent of $\theta \in \angle(\alpha, \beta)$.

Corollary. If there is a sector $\angle(\alpha', \beta') \ni \angle(\alpha, \beta)$ such that $\alpha' < \alpha < \beta < \beta'$, and the weight M from Theorem 4 is radial and differentiable in r , then, for a from small neighborhood of 1, for all assertions (Z), (U) и (M) of this Theorem at points $z \in \angle(\alpha, \beta)$ the summand $\frac{C_\varepsilon}{2-a} q_M^{[a]}(z)$ in (17) can be change to a summand

$$\frac{aC_\varepsilon}{2(2-a)} \frac{1}{\sqrt{1-|z|}} \int_{(|z|-a\sqrt{1-|z|^2})^+}^1 (1-t) d(tM'(t)). \quad (20)$$

for small $a > 1$.

This work is supported by the RFSB grants No. 09-01-00046-a, No. 08-01-97023-Volga region, and by the grant of President of Russian Federation “State support of leading scientific schools”, project 3081.2008.1.

REFERENCES

- [1] Andriy Kondratyuk, and Ilpo Laine, Meromorphic functions in multiply connected domains, Joensuu–L’viv, 2006, pp. 9–111
- [2] Khabibullin B. N. *Zero sequences of holomorphic functions, the representation of meromorphic functions and harmonic minorants* // Matematicheskii Sbornik – 2007. – **198**:2. – 121–160 (in Russian). Eng. transl. in “Sbornik: Mathematics”. 2007. – **198**:2. – 261–298.
- [3] Khabibullin B. N. *Dual representation of superlinear functionals and its applications in function theory. II* // Izvestiya RAN – 2001. – **65**:5. – 167–190; Eng. transl. in “Izvestiya: Mathematics” – 2001. – **65**:5. – 1017–1039.
- [4] Kudasheva E. G., and Khabibullin B. N. *The distribution of the zeros of holomorphic functions of moderate growth in the unit disc and the representation of meromorphic functions* // Matematicheskii Sbornik – 2009. – **200**:9 – 95–126 (in Russian); Eng. transl. in “Sbornik: Mathematics” – 2009. – **198**:2 – 1353–1382.

Russia, Ufa, Bashkir State University, khabib-bulat@mail.ru